

# Interactions with Experimental COVID-19 Adjunct Therapies

Charts updated 5 October 2021

Page 1 of 6

Please check [www.covid19-druginteractions.org](http://www.covid19-druginteractions.org) for updates.

Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Drug interaction data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

	Aspirin	Dalteparin	Enoxaparin
<b>Anaesthetics &amp; Muscle Relaxants</b>			
Alcuronium	◆	◆	◆
Bupivacaine	◆	◆	◆
Cisatracurium	◆	◆	◆
Desflurane	◆	◆	◆
Dexmedetomidine	◆	◆	◆
Enflurane	◆	◆	◆
Ephedrine	◆	◆	◆
Etidocaine	◆	◆	◆
Halothane	◆	◆	◆
Isoflurane	◆	◆	◆
Ketamine	◆	◆	◆
Minaxolone	◆	◆	◆
Nitrous oxide	◆	◆	◆
Propofol	◆	◆	◆
Rocuronium	◆	◆	◆
Sevoflurane	◆	◆	◆
Sufentanil	◆	◆	◆
Suxamethonium (succinylcholine)	◆	◆	◆
Tetracaine	◆	◆	◆
Thiopental	◆	◆	◆
Tizanidine	◆	◆	◆
Vecuronium	◆	◆	◆
<b>Analgesics</b>			
Alfentanil	◆	◆	◆
Aspirin	◆	■	■
Buprenorphine	◆	◆	◆
Celecoxib	◆	■	■
Codeine	◆	◆	◆
Dextropropoxyphene	◆	◆	◆
Diamorphine (diacetylmorphine)	◆	◆	◆
Diclofenac	◆	■	■
Dihydrocodeine	◆	◆	◆
Fentanyl	◆	◆	◆
Hydrocodone	◆	◆	◆
Hydromorphone	◆	◆	◆
Ibuprofen	◆	■	■
Mefenamic acid	◆	■	■
Metamizole	■	◆	◆
Methadone	◆	◆	◆
Morphine	◆	◆	◆
Naproxen	◆	■	■
Nimesulide	◆	◆	◆
Oxycodone	◆	◆	◆
Paracetamol (Acetaminophen)	◆	◆	◆
Pethidine (Meperidine)	◆	◆	◆
Piroxicam	◆	■	■
Remifentanyl	◆	◆	◆
Tapentadol	◆	◆	◆
Tramadol	◆	◆	◆
<b>Antiarrhythmics</b>			
Amiodarone	◆	◆	◆
Bepidil	◆	◆	◆
Digoxin	◆	■	◆
Disopyramide	◆	◆	◆
Dofetilide	◆	◆	◆
Flecainide	◆	◆	◆
Lidocaine (Lignocaine)	◆	◆	◆
Mexiletine	◆	◆	◆
Propafenone	◆	◆	◆
Quinidine	◆	◆	◆

	Aspirin	Dalteparin	Enoxaparin
<b>Antibacterials</b>			
Amikacin	◆	◆	◆
Amoxicillin	◆	◆	◆
Ampicillin	◆	◆	◆
Bedaquiline	◆	◆	◆
Capreomycin	◆	◆	◆
Cefalexin	◆	◆	◆
Cefazolin	◆	◆	◆
Cefepime	◆	◆	◆
Cefixime	◆	◆	◆
Cefotaxime	◆	◆	◆
Ceftazidime	◆	◆	◆
Ceftriaxone	◆	◆	◆
Chloramphenicol	◆	◆	◆
Ciprofloxacin	◆	◆	◆
Clarithromycin	◆	◆	◆
Clavulanic acid	◆	◆	◆
Clindamycin	◆	◆	◆
Clofazimine	◆	◆	◆
Cloxacillin	◆	◆	◆
Cycloserine	◆	◆	◆
Dapsone	◆	◆	◆
Delamanid	◆	◆	◆
Doxycycline	◆	◆	◆
Ertapenem	◆	◆	◆
Erythromycin	◆	◆	◆
Ethambutol	◆	◆	◆
Ethionamide	◆	◆	◆
Flucloxacillin	◆	◆	◆
Gentamicin	◆	◆	◆
Imipenem/Cilastatin	◆	◆	◆
Isoniazid	◆	◆	◆
Kanamycin	◆	◆	◆
Levofloxacin	◆	◆	◆
Linezolid	◆	◆	◆
Meropenem	◆	◆	◆
Metronidazole	◆	◆	◆
Moxifloxacin	◆	◆	◆
Nitrofurantoin	◆	◆	◆
Ofloxacin	◆	◆	◆
Para-aminosalicylic acid	◆	◆	◆
Penicillins	◆	▲	◆
Piperacillin	◆	◆	◆
Pyrazinamide	◆	◆	◆
Rifabutin	◆	◆	◆
Rifampicin	◆	◆	◆
Rifapentine	◆	◆	◆
Rifaximin	◆	◆	◆
Spectinomycin	◆	◆	◆
Streptomycin	◆	◆	◆
Sulfadiazine	◆	◆	◆
Tazobactam	◆	◆	◆
Telithromycin	◆	◆	◆
Tetracyclines	◆	■	◆
Tinidazole	◆	◆	◆
Trimethoprim/Sulfamethoxazole	◆	◆	◆
Vancomycin	◆	◆	◆

## Key to symbols

●	These drugs should not be coadministered
■	Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

© Liverpool Drug Interaction Group, University of Liverpool, Pharmacology Research Labs, 1st Floor Block H, 70 Pembroke Place, LIVERPOOL, L69 3GF

We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.

# Interactions with Experimental COVID-19 Adjunct Therapies

Charts updated 5 October 2021

Page 2 of 6

Please check [www.covid19-druginteractions.org](http://www.covid19-druginteractions.org) for updates.

Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Drug interaction data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

	Aspirin	Dalteparin	Enoxaparin
<b>Anti-coagulant, Anti-platelet and Fibrinolytic</b>			
Acenocoumarol	■	■	■
Apixaban	■	■	■
Argatroban	■	■	■
Aspirin (anti-platelet)	■	■	■
Betrixaban	■	■	■
Clopidogrel	■	■	■
Dabigatran	■	■	■
Dalteparin	■	■	■
Dipyridamole	■	■	■
Edoxaban	■	■	■
Enoxaparin	■	■	■
Fondaparinux	■	■	■
Heparin	■	■	■
Phenprocoumon	■	■	■
Prasugrel	■	■	■
Rivaroxaban	■	■	■
Streptokinase	■	■	■
Ticagrelor	■	■	■
Tinzaparin	■	■	■
Warfarin	■	■	■
<b>Anticonvulsants</b>			
Brivaracetam	◆	◆	◆
Carbamazepine	◆	◆	◆
Clonazepam	◆	◆	◆
Eslicarbazepine	◆	◆	◆
Ethosuximide	◆	◆	◆
Gabapentin	◆	◆	◆
Lacosamide	◆	◆	◆
Lamotrigine	◆	◆	◆
Levetiracetam	◆	◆	◆
Oxcarbazepine	◆	◆	◆
Perampanel	◆	◆	◆
Phenobarbital (Phenobarbitone)	◆	◆	◆
Phenytoin	▲	◆	◆
Pregabalin	◆	◆	◆
Primidone	◆	◆	◆
Retigabine	◆	◆	◆
Rufinamide	◆	◆	◆
Sultiame	◆	◆	◆
Tiagabine	◆	◆	◆
Topiramate	◆	◆	◆
Valproate (Divalproex)	▲	◆	◆
Vigabatrin	◆	◆	◆
Zonisamide	◆	◆	◆

	Aspirin	Dalteparin	Enoxaparin
<b>Antidepressants</b>			
Agomelatine	◆	◆	◆
Amitriptyline	◆	◆	◆
Bupropion	◆	◆	◆
Citalopram	◆	◆	◆
Clomipramine	◆	◆	◆
Desipramine	◆	◆	◆
Doxepin	◆	◆	◆
Duloxetine	◆	◆	◆
Escitalopram	◆	◆	◆
Fluoxetine	◆	◆	◆
Fluvoxamine	◆	◆	◆
Imipramine	◆	◆	◆
Lithium	◆	◆	◆
Maprotiline	◆	◆	◆
Mianserin	◆	◆	◆
Milnacipran	◆	◆	◆
Mirtazapine	◆	◆	◆
Nefazodone	◆	◆	◆
Nortriptyline	◆	◆	◆
Paroxetine	◆	◆	◆
Phenelzine	◆	◆	◆
Reboxetine	◆	◆	◆
Sertraline	◆	◆	◆
St John's wort	◆	◆	◆
Tranlycypromine	◆	◆	◆
Trazodone	◆	◆	◆
Trimipramine	◆	◆	◆
Venlafaxine	◆	◆	◆
Vortioxetine	◆	◆	◆
<b>Anti-diabetics</b>			
Acarbose	◆	◆	◆
Canagliflozin	◆	◆	◆
Dapagliflozin	◆	◆	◆
Dulaglutide	◆	◆	◆
Empagliflozin	◆	◆	◆
Exenatide	◆	◆	◆
Glibenclamide (Glyburide)	◆	◆	◆
Gliclazide	◆	◆	◆
Glimepiride	◆	◆	◆
Glipizide	◆	◆	◆
Insulin	◆	◆	◆
Linagliptin	◆	◆	◆
Liraglutide	◆	◆	◆
Metformin	◆	◆	◆
Nateglinide	◆	◆	◆
Pioglitazone	◆	◆	◆
Repaglinide	◆	◆	◆
Rosiglitazone	◆	◆	◆
Saxagliptin	◆	◆	◆
Sitagliptin	◆	◆	◆
Tolbutamide	◆	◆	◆
Vildagliptin	◆	◆	◆

### Key to symbols

●	These drugs should not be coadministered
■	Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

## Interactions with Experimental COVID-19 Adjunct Therapies

Charts updated 5 October 2021

Page 3 of 6

Please check [www.covid19-druginteractions.org](http://www.covid19-druginteractions.org) for updates.

Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Drug interaction data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

	Aspirin	Dalteparin	Enoxaparin
<b>Antifungals</b>			
Amphotericin B	◆	◆	◆
Anidulafungin	◆	◆	◆
Caspofungin	◆	◆	◆
Fluconazole	◆	◆	◆
Flucytosine	◆	◆	◆
Griseofulvin	◆	◆	◆
Isavuconazole	◆	◆	◆
Itraconazole	◆	◆	◆
Ketoconazole	◆	◆	◆
Micafungin	◆	◆	◆
Miconazole	◆	◆	◆
Nystatin	◆	◆	◆
Posaconazole	◆	◆	◆
Terbinafine	◆	◆	◆
Voriconazole	◆	◆	◆
<b>Antipsychotics/Neuroleptics</b>			
Amisulpride	◆	◆	◆
Aripiprazole	◆	◆	◆
Asenapine	◆	◆	◆
Chlorpromazine	◆	◆	◆
Clozapine	◆	◆	◆
Fluphenazine	◆	◆	◆
Haloperidol	◆	◆	◆
Iloperidone	◆	◆	◆
Levomepromazine	◆	◆	◆
Olanzapine	◆	◆	◆
Paliperidone	◆	◆	◆
Perazine	◆	◆	◆
Periciazine	◆	◆	◆
Perphenazine	◆	◆	◆
Pimozide	◆	◆	◆
Pipotiazine	◆	◆	◆
Quetiapine	◆	◆	◆
Risperidone	◆	◆	◆
Sulpiride	◆	◆	◆
Thioridazine	◆	◆	◆
Tiapride	◆	◆	◆
Ziprasidone	◆	◆	◆
Zotepine	◆	◆	◆
Zuclopenthixol	◆	◆	◆

	Aspirin	Dalteparin	Enoxaparin
<b>Anxiolytics/Hypnotics/Sedatives</b>			
Alprazolam	◆	◆	◆
Bromazepam	◆	◆	◆
Buspirone	◆	◆	◆
Chlordiazepoxide	◆	◆	◆
Clobazam	◆	◆	◆
Clorazepate	◆	◆	◆
Diazepam	◆	◆	◆
Estazolam	◆	◆	◆
Flunitrazepam	◆	◆	◆
Flurazepam	◆	◆	◆
Hydroxyzine	◆	◆	◆
Lorazepam	◆	◆	◆
Lormetazepam	◆	◆	◆
Midazolam (oral)	◆	◆	◆
Midazolam (parenteral)	◆	◆	◆
Oxazepam	◆	◆	◆
Temazepam	◆	◆	◆
Triazolam	◆	◆	◆
Zaleplon	◆	◆	◆
Zolpidem	◆	◆	◆
Zopiclone	◆	◆	◆
<b>Beta Blockers</b>			
Atenolol	◆	◆	◆
Bisoprolol	◆	◆	◆
Carvedilol	◆	◆	◆
Metoprolol	◆	◆	◆
Nebivolol	◆	◆	◆
Oxprenolol	◆	◆	◆
Pindolol	◆	◆	◆
Propranolol	◆	◆	◆
Timolol	◆	◆	◆
<b>Bronchodilators</b>			
Acidinium bromide	◆	◆	◆
Aminophylline	◆	◆	◆
Formoterol	◆	◆	◆
Glycopyrronium bromide	◆	◆	◆
Indacaterol	◆	◆	◆
Ipratropium bromide	◆	◆	◆
Montelukast	◆	◆	◆
Olodaterol	◆	◆	◆
Roflumilast	◆	◆	◆
Salbutamol	◆	◆	◆
Salmeterol	◆	◆	◆
Theophylline	◆	◆	◆
Tiotropium bromide	◆	◆	◆
Umeclidinium bromide	◆	◆	◆
Vilanterol	◆	◆	◆
<b>Calcium Channel Blockers</b>			
Amlodipine	◆	◆	◆
Diltiazem	◆	◆	◆
Felodipine	◆	◆	◆
Nicardipine	◆	◆	◆
Nifedipine	◆	◆	◆
Nisoldipine	◆	◆	◆
Nitrendipine	◆	◆	◆
Verapamil	◆	◆	◆

## Key to symbols

●	These drugs should not be coadministered
■	Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

© Liverpool Drug Interaction Group, University of Liverpool, Pharmacology Research Labs, 1st Floor Block H, 70 Pembroke Place, LIVERPOOL, L69 3GF

We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.

# Interactions with Experimental COVID-19 Adjunct Therapies

Charts updated 5 October 2021

Page 4 of 6

Please check [www.covid19-druginteractions.org](http://www.covid19-druginteractions.org) for updates.

Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Drug interaction data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

	Aspirin	Dalteparin	Enoxaparin
<b>Contraceptives/HRT</b>			
Desogestrel (COC)	◆	◆	◆
Desogestrel (POP)	◆	◆	◆
Drospirenone (COC)	◆	◆	◆
Drospirenone (HRT)	◆	◆	◆
Dydrogesterone (HRT)	◆	◆	◆
Estradiol	◆	◆	◆
Ethinylestradiol	◆	◆	◆
Etonogestrel (implant)	◆	◆	◆
Etonogestrel (vaginal ring)	◆	◆	◆
Gestodene (COC)	◆	◆	◆
Levonorgestrel (COC)	◆	◆	◆
Levonorgestrel (emergency contrac.)	◆	◆	◆
Levonorgestrel (HRT)	◆	◆	◆
Levonorgestrel (implant)	◆	◆	◆
Levonorgestrel (IUD)	◆	◆	◆
Levonorgestrel (POP)	◆	◆	◆
Medroxyprogesterone (depot inj.)	◆	◆	◆
Medroxyprogesterone (oral)	◆	◆	◆
Norelgestromin (patch)	◆	◆	◆
Norethisterone (COC)	◆	◆	◆
Norethisterone (HRT)	◆	◆	◆
Norethisterone (IM depot)	◆	◆	◆
Norethisterone (POP)	◆	◆	◆
Norgestimate (COC)	◆	◆	◆
Norgestrel (COC)	◆	◆	◆
Norgestrel (HRT)	◆	◆	◆
Ulipristal	◆	◆	◆
<b>Covid-19 Adjunct Therapies</b>			
Aspirin (Covid-19 adjunct)	■	■	■
Dalteparin (Covid-19 adjunct)	■	■	■
Enoxaparin (Covid 19 adjunct)	■	■	■
<b>Covid-19 Antiviral Therapies</b>			
Atazanavir	◆	◆	◆
Azithromycin	◆	◆	◆
Bamlanivimab/Etesevimab	◆	◆	◆
Casirivimab/Imdevimab	◆	◆	◆
Chloroquine	◆	◆	◆
Favipiravir	◆	◆	◆
Hydroxychloroquine	◆	◆	◆
Interferon beta	◆	◆	◆
Ivermectin	◆	◆	◆
Lopinavir/ritonavir	◆	◆	◆
Molnupiravir	◆	◆	◆
Niclosamide	◆	◆	◆
Nitazoxanide	◆	◆	◆
Remdesivir	◆	◆	◆
Ribavirin	◆	◆	◆
<b>Covid-19 Immune Therapies</b>			
Anakinra	◆	◆	◆
Baricitinib	◆	◆	◆
Budesonide (inhaled)	◆	◆	◆
Canakinumab	◆	◆	◆
Colchicine	◆	◆	◆
Covid 19 convalescent plasma	◆	◆	◆
Covid-19 vaccines	◆	◆	◆
Dexamethasone (low dose)	■	◆	◆
Hydrocortisone (oral or IV)	■	◆	◆
Methylprednisolone (oral or IV)	■	◆	◆
Ruxolitinib	◆	◆	◆
Sarilumab	◆	◆	◆
Tocilizumab	◆	◆	◆

	Aspirin	Dalteparin	Enoxaparin
<b>Gastrointestinal Agents</b>			
Alosetron	◆	◆	◆
Antacids	◆	◆	◆
Bisacodyl	◆	◆	◆
Cimetidine	◆	◆	◆
Cisapride	◆	◆	◆
Esomeprazole	◆	◆	◆
Famotidine	◆	◆	◆
Lactulose	◆	◆	◆
Lansoprazole	◆	◆	◆
Loperamide	◆	◆	◆
Magnesium salts (oral)	◆	◆	◆
Mesalazine	◆	◆	◆
Omeprazole	◆	◆	◆
Pantoprazole	◆	◆	◆
Prucalopride	◆	◆	◆
Rabeprazole	◆	◆	◆
Ranitidine	◆	◆	◆
Senna	◆	◆	◆
<b>Gastrointestinal Agents (anti-emetics)</b>			
Aprepitant	◆	◆	◆
Cyclizine	◆	◆	◆
Dolasetron	◆	◆	◆
Domperidone	◆	◆	◆
Dronabinol	◆	◆	◆
Granisetron	◆	◆	◆
Metoclopramide	◆	◆	◆
Ondansetron	◆	◆	◆
Prochlorperazine	◆	◆	◆
<b>HCV DAAs</b>			
Elbasvir/Grazoprevir	◆	◆	◆
Glecaprevir/Pibrentasvir	◆	◆	◆
Ledipasvir/Sofosbuvir	◆	◆	◆
Ombitasvir/Paritaprevir/r	◆	◆	◆
Ombitasvir/Paritaprevir/r + Dasabuvir	◆	◆	◆
Sofosbuvir	◆	◆	◆
Sofosbuvir/Velpatasvir	◆	◆	◆
Sofosbuvir/Velpatasvir/Voxilaprevir	◆	◆	◆

### Key to symbols

●	These drugs should not be coadministered
■	Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

# Interactions with Experimental COVID-19 Adjunct Therapies

Charts updated 5 October 2021

Page 5 of 6

Please check [www.covid19-druginteractions.org](http://www.covid19-druginteractions.org) for updates.

Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Drug interaction data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

	Aspirin	Dalteparin	Enoxaparin
<b>HIV Antiretroviral Therapies</b>			
Abacavir	◆	◆	◆
Albuvirtide	◆	◆	◆
Atazanavir + ritonavir	◆	◆	◆
Atazanavir/cobicistat	◆	◆	◆
Bictegravir/Emtricitabine/TAF	◆	◆	◆
Cabotegravir (oral)	◆	◆	◆
Cabotegravir/rilpivirine (long acting)	◆	◆	◆
Darunavir + ritonavir	◆	◆	◆
Darunavir/cobicistat	◆	◆	◆
Darunavir/Cobi/Emtricitabine/TAF	◆	◆	◆
Dolutegravir	◆	◆	◆
Dolutegravir/Lamivudine	◆	◆	◆
Dolutegravir/Rilpivirine	◆	◆	◆
Dolutegravir/Abacavir/Lamivudine	◆	◆	◆
Doravirine	◆	◆	◆
Doravirine/Lamivudine/Tenofovir-DF	◆	◆	◆
Efavirenz	◆	◆	◆
Elvitegravir/Cobi/Emtricitabine/TAF	◆	◆	◆
Elvitegravir/Cobi/Emtricitabine/TDF	◆	◆	◆
Emtricitabine	◆	◆	◆
Emtricitabine/Tenofovir alafenamide	◆	◆	◆
Emtricitabine/Tenofovir-DF	◆	◆	◆
Etravirine	◆	◆	◆
Fostemsavir	◆	◆	◆
Ibalizumab-uiyk	◆	◆	◆
Lamivudine	◆	◆	◆
Maraviroc	◆	◆	◆
Nevirapine	◆	◆	◆
Raltegravir	◆	◆	◆
Rilpivirine	◆	◆	◆
Rilpivirine/Emtricitabine/TAF	◆	◆	◆
Tenofovir-DF	◆	◆	◆
Zidovudine	◆	◆	◆

	Aspirin	Dalteparin	Enoxaparin
<b>Hypertension / Heart Failure Agents</b>			
Aliskiren	◆	◆	◆
Ambrisentan	◆	◆	◆
Amiloride	◆	■	■
Benazepril	◆	◆	◆
Bendroflumethiazide	◆	◆	◆
Bosentan	◆	◆	◆
Candesartan	◆	◆	◆
Captopril	◆	◆	◆
Chlortalidone	◆	◆	◆
Cilazapril	◆	◆	◆
Clonidine	◆	◆	◆
Dopamine	◆	◆	◆
Doxazosin	◆	◆	◆
Enalapril	◆	◆	◆
Eplerenone	◆	■	■
Epoprostenol	■	■	■
Eprosartan	◆	◆	◆
Fosinopril	◆	◆	◆
Furosemide	◆	◆	◆
Hydralazine	◆	◆	◆
Hydrochlorothiazide	◆	◆	◆
Iloprost	■	■	■
Indapamide	◆	◆	◆
Irbesartan	◆	◆	◆
Isosorbide dinitrate	◆	◆	◆
Ivabradine	◆	◆	◆
Labetalol	◆	◆	◆
Lacidipine	◆	◆	◆
Lercanidipine	◆	◆	◆
Lisinopril	◆	◆	◆
Losartan	◆	◆	◆
Macitentan	◆	◆	◆
Methyldopa	◆	◆	◆
Metolazone	◆	◆	◆
Moxonidine	◆	◆	◆
Olmesartan	◆	◆	◆
Perindopril	◆	◆	◆
Prazosin	◆	◆	◆
Quinapril	◆	◆	◆
Ramipril	◆	◆	◆
Ranolazine	◆	◆	◆
Riociguat	◆	◆	◆
Sacubitril	◆	◆	◆
Selexipag	◆	◆	◆
Sildenafil (PAH)	◆	◆	◆
Sodium nitroprusside	◆	◆	◆
Spirolactone	◆	■	■
Tadalafil (PAH)	◆	◆	◆
Telmisartan	◆	◆	◆
Terazosin	◆	◆	◆
Torsemide	◆	◆	◆
Trandolapril	◆	◆	◆
Treprostinil	■	■	■
Valsartan	◆	◆	◆
Xipamide	◆	◆	◆

### Key to symbols

●	These drugs should not be coadministered
■	Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

© Liverpool Drug Interaction Group, University of Liverpool, Pharmacology Research Labs, 1st Floor Block H, 70 Pembroke Place, LIVERPOOL, L69 3GF

We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.

## Interactions with Experimental COVID-19 Adjunct Therapies

Charts updated 5 October 2021

Page 6 of 6

Please check [www.covid19-druginteractions.org](http://www.covid19-druginteractions.org) for updates.

Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Drug interaction data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

	Aspirin	Dalteparin	Enoxaparin
<b>Immunosuppressants</b>			
Adalimumab	◆	◆	◆
Anti-thymocyte globulin	◆	◆	◆
Azathioprine	◆	◆	◆
Basiliximab	◆	◆	◆
Belatacept	◆	◆	◆
Ciclosporin	◆	◆	◆
Everolimus	◆	◆	◆
Methotrexate	◆	◆	◆
Mycophenolate	◆	◆	◆
Pirfenidone	◆	◆	◆
Sirolimus	◆	◆	◆
Tacrolimus	◆	◆	◆
<b>Inotropes &amp; Vasopressors</b>			
Adrenaline (Epinephrine)	◆	◆	◆
Dobutamine	◆	◆	◆
Noradrenaline	◆	◆	◆
Vasopressin	◆	◆	◆
<b>Lipid Lowering Agents</b>			
Atorvastatin	◆	◆	◆
Bezafibrate	◆	◆	◆
Clofibrate	◆	◆	◆
Evolocumab	◆	◆	◆
Ezetimibe	◆	◆	◆
Fenofibrate	◆	◆	◆
Fish oils	◆	◆	◆
Fluvastatin	◆	◆	◆
Gemfibrozil	◆	◆	◆
Lovastatin	◆	◆	◆
Pitavastatin	◆	◆	◆
Pravastatin	◆	◆	◆
Rosuvastatin	◆	◆	◆
Simvastatin	◆	◆	◆
<b>Other</b>			
Acetylcysteine	◆	◆	◆
Aciclovir	◆	◆	◆
Alendronic acid	◆	◆	◆
Alfuzosin	◆	◆	◆
Allopurinol	◆	◆	◆
Calcium supplements	◆	◆	◆
Carbocysteine	◆	◆	◆
Dextromethorphan	◆	◆	◆
Donepezil	◆	◆	◆
Eltrombopag	◆	◆	◆
Entecavir	◆	◆	◆
Finasteride	◆	◆	◆
Folic acid	◆	◆	◆
Guaifenesin	◆	◆	◆
Levothyroxine	◆	◆	◆
Magnesium sulphate (IV)	◆	◆	◆
Melatonin	◆	◆	◆
Memantine	◆	◆	◆
Mirabegron	◆	◆	◆
Potassium	◆	◆	◆
Pramipexole	◆	◆	◆
Pyridostigmine	◆	◆	◆
Oseltamivir	◆	◆	◆
Tamsulosin	◆	◆	◆

	Aspirin	Dalteparin	Enoxaparin
<b>Steroids</b>			
Beclometasone	■	◆	◆
Betamethasone	■	◆	◆
Budesonide (oral/rectal)	■	◆	◆
Ciclesonide	◆	◆	◆
Clobetasol	◆	◆	◆
Fludrocortisone	■	◆	◆
Flunisolide	◆	◆	◆
Fluocinolone	◆	◆	◆
Fluticasone	◆	◆	◆
Hydrocortisone (topical)	◆	◆	◆
Megestrol acetate	◆	◆	◆
Methylprednisolone (topical)	◆	◆	◆
Mometasone	◆	◆	◆
Nandrolone	◆	◆	◆
Oxandrolone	◆	◆	◆
Prednisolone	■	◆	◆
Prednisone	■	◆	◆
Stanozolol	◆	◆	◆
Testosterone	◆	◆	◆
Triamcinolone	■	◆	◆

## Key to symbols

●	These drugs should not be coadministered
■	Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

© Liverpool Drug Interaction Group, University of Liverpool, Pharmacology Research Labs, 1st Floor Block H, 70 Pembroke Place, LIVERPOOL, L69 3GF

We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.