

COVID-19 Therapies - Dose Recommendations for Patients with Renal Impairment

Charts updated 18 March 2022

Page 1 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

COVID-19 Antiviral Therapies (Licensed or Under Clinical Investigation)

Please refer to the additional notes on the following pages for further details and complete dose recommendations.

Drug	Renal Impairment eGFR*				Renal Replacement Therapy (RRT)			References
	≥50 ml/min	30-49 ml/min	10-29 ml/min	<10 ml/min	Haemodialysis	CRRT	PD	
Azithromycin	100%	100%	100%	100%, with caution	100%, with caution	100%, with caution	100%, with caution	1-4
Bamlanivimab/ Etesevimab	100%	100%	100%	100%	100%	100%	100%	5
Casirivimab/ Imdevimab	100%	100%	100%	100%	100%	100%	100%	6, 7
Favipiravir	100%	No recommendation possible	No recommendation possible	No recommendation possible	No recommendation possible	No recommendation possible	No recommendation possible	8
Molnupiravir	100%	100%	100%	100%	100%	100%	100%	9, 10
Niclosamide	100%	100%	100%	100%	100%	100%	100%	11, 12
Nirmatrelvir + ritonavir	100%	Nirmatrelvir: 50% Ritonavir: 100% (i.e. 1 tablet of each)	Not recommended	Not recommended	Not recommended	Not recommended	Not recommended	13, 14
Nitazoxanide	100%	100%, with caution	100%, with caution	100%, with caution	100%, with caution	100%, with caution	100%, with caution	15
Remdesivir	100%	100%	Not recommended	Not recommended	Not recommended	Not recommended	Not recommended	16, 17
Sotrovimab	100%	100%	100%	100%	100%	100%	100%	18, 19
Tixagevimab/ Cilgavimab	100%	100%	100%	100%	100%	100%	100%	20

Abbreviations

eGFR Estimated glomerular filtration rate

* Use CKD-EPI formula: the Abbreviated Modification of Diet in Renal Disease (aMDRD) or the Cockcroft-Gault (CG) equation may be used as an alternative; see www.chip.dk/Tools-Standards/Clinical-risk-scores

CRRT Continuous renal replacement therapies

PD Peritoneal dialysis

COVID-19 Therapies - Dose Recommendations for Patients with Renal Impairment

Charts updated 18 March 2022

Page 2 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

Notes - Antiviral Therapies (Licensed or Under Clinical Investigation)

Azithromycin

Approximately 6-12% of an IV dose of azithromycin is excreted unchanged in urine.

In patients with severe renal impairment (GFR <10 ml/min) systemic exposure to azithromycin increased by 33-35%.

The extent to which azithromycin is removed with haemodialysis or CVVH is unknown; use with caution. Azithromycin is not substantially removed by CAPD.

Bamlanivimab/ Etesevimab

Bamlanivimab and etesevimab are not eliminated intact in the urine. Renal impairment is not expected to affect the exposure of bamlanivimab or etesevimab.

No dosage adjustment is recommended in patients with renal impairment.

Casirivimab/ Imdevimab

Casirivimab and imdevimab are monoclonal antibodies and are therefore not likely to undergo renal excretion. Renal impairment is not expected to affect the exposure of casirivimab and imdevimab

No dosage adjustment is required in individuals with mild or moderate renal impairment, or in patients with creatinine clearance (CrCl) < 15 mL/min including those on dialysis.

Based on population PK analysis, trough concentrations of casirivimab and imdevimab in serum at steady state were comparable between patients with mild or moderate renal impairment, or patients with CrCl <15 ml/min including those on dialysis, and patients with normal renal function. Limited data are available in patients with severe renal impairment (n=3).

Favipiravir

Favipiravir is 90.5% renally excreted, the majority of which (82-92%) as M1 metabolite which is responsible for toxicity.

M1 may accumulate in renal impairment with a 2.5-fold increase in moderate impairment based on a single patient studied in global phase 3 with eGFR

30-50 ml/min. Uric acid increases may also be a concern in renal impairment. No data is available to make any statement of safety in patients with renal impairment or dependent on RRT.

Molnupiravir

Molnupiravir is hydrolysed to n-hydroxycytidine (NHC) prior to reaching systemic circulation. Renal clearance is not a meaningful route of elimination for NHC.

No dose adjustment in patients with any degree of renal impairment is needed. In a population PK analysis, mild or moderate renal impairment did not have a meaningful impact on the pharmacokinetics of NHC. The pharmacokinetics of molnupiravir and NHC has not been evaluated in patients with eGFR <30 mL/min or on dialysis.

Niclosamide

Niclosamide has poor bioavailability when administered orally and low systemic absorption via the intranasal/inhaled route.

Niclosamide may be given safely to patients with kidney diseases.

Nirmatrelvir +ritonavir

The primary route of elimination of nirmatrelvir when administered with ritonavir was renal excretion of intact drug. Approximately 49.6% of the administered dose was recovered in urine.

Compared to healthy controls with no renal impairment, nirmatrelvir C_{max} and AUC were 30% and 24% higher in patients with mild renal impairment, 38% and 87% higher in patients with moderate renal impairment, and 48% and 204% higher in patients with severe renal impairment.

Nirmatrelvir and ritonavir are highly protein bound, therefore, it is unlikely that they will be significantly removed by haemodialysis or peritoneal dialysis.

COVID-19 Therapies - Dose Recommendations for Patients with Renal Impairment

Charts updated 18 March 2022

Page 3 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

Notes - Antiviral Therapies (Licensed or Under Clinical Investigation) [continued]

Nitazoxanide

Approximately one third of an oral dose of nitazoxanide is excreted in the urine.

Nitazoxanide has not been studied in patients with compromised renal function. No data are available in RRT.

Remdesivir

Remdesivir is not cleared unchanged in urine to any substantial extent, but its main metabolite GS-441524 is renally cleared and the metabolite levels in plasma may theoretically increase in patients with impaired renal function. The excipient betadex sulfobutyl ether sodium is renally cleared and accumulates in patients with decreased renal function.

The pharmacokinetics of remdesivir have not been evaluated in patients with renal impairment. Patients with eGFR ≥ 30 mL/min have received remdesivir for treatment of COVID-19 with no dose adjustment.

All patients should have eGFR determined prior to starting remdesivir and while receiving it as clinically appropriate. Remdesivir should not be used in patients with eGFR < 30 mL/min as administration of drugs formulated with betadex sulfobutyl ether sodium (such as remdesivir) is not recommended in patients with eGFR < 30 mL/min.

Note, a renal toxicity signal has been noticed for remdesivir and is currently under investigation by the European Medicines Agency (Gérard AO *et al.* [Clin Pharmacol Ther, 2021, 109\(4\):1021-1024](#)).

Sotrovimab

Renal impairment is not expected to impact the pharmacokinetics of sotrovimab since mAbs with molecular weight > 69 kDa do not undergo renal elimination.

No dosage adjustment is recommended in patients with renal impairment.

Due to its molecular weight, dialysis is not expected to impact the pharmacokinetics of sotrovimab.

Tixagevimab/ Cilgavimab

Tixagevimab and cilgavimab are monoclonal antibodies and are therefore not likely to undergo renal excretion. Renal impairment is not expected to affect the exposure of tixagevimab and cilgavimab

There is no difference in the clearance of tixagevimab and cilgavimab in individuals with mild or moderate renal impairment compared to individuals with normal renal function. There were insufficient subjects with severe renal impairment to draw conclusions.

COVID-19 Therapies - Dose Recommendations for Patients with Renal Impairment

Charts updated 18 March 2022

Page 4 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

COVID-19 Host-directed Therapies (Licensed or Under Clinical Investigation)

Please refer to the additional notes on the following pages for further details and complete dose recommendations.

Drug	Renal Impairment eGFR*				Renal Replacement Therapy (RRT)			References
	≥50 ml/min	30-49 ml/min	10-29 ml/min	<10 ml/min	Haemodialysis	CRRT	PD	
Anakinra	100%	100%	100% every other day	100% every other day	100% every other day	100% every other day	100% every other day	21, 22
Baricitinib	100%	2 mg once daily	Not recommended	Not recommended	Not recommended	2 mg once daily	Not recommended	4, 23, 24
Budesonide (inhaled)	100%	100%	100%	100%	100%	100%	100%	4, 25, 26
Canakinumab	100%	100%	100%	100%	100%	100%	100%	27, 28
Dexamethasone (low dose)	100%	100%	100%	100%	100%	100%	100%	4, 29
Fluvoxamine	100%	100%	100%	100% following slow titration	100% following slow titration	100% following slow titration	100% following slow titration	4, 30, 31
Hydrocortisone	100%	100%, with caution	100%, with caution	100%, with caution	100%	100%	100%	4, 32
Infliximab	100%	100%, with caution	100%, with caution	100%, with caution	100%, with caution	100%, with caution	100%, with caution	4, 33, 34
Methylprednisolone	100%	100%	100%	100%	100%	100%	100%	4, 35, 36
Ruxolitinib	100%	100%	5 mg twice daily	Not recommended	10 mg single dose or 5 mg twice daily	Use with caution. Consider dosing as for eGFR 10-30 ml/min	Use with caution. Consider dosing as for eGFR 10-30 ml/min	4, 37-39
Sarilumab	100%	100%	100%	100%	100%	100%	100%	40
Tocilizumab	100%	100%	100%	100%	100%	100%	100%	4, 41, 42

Abbreviations

eGFR Estimated glomerular filtration rate

* Use CKD-EPI formula: the Abbreviated Modification of Diet in Renal Disease (aMDRD) or the Cockcroft-Gault (CG) equation may be used as an alternative; see www.chip.dk/Tools-Standards/Clinical-risk-scores

CRRT Continuous renal replacement therapies

PD Peritoneal dialysis

COVID-19 Therapies - Dose Recommendations for Patients with Renal Impairment

Charts updated 18 March 2022

Page 5 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

Notes – Host-directed Therapies (Licensed or Under Clinical Investigation)

- Anakinra** *Anakinra is eliminated by glomerular filtration and subsequent tubular metabolism. Plasma clearance of anakinra decreases with decreasing renal function. Plasma clearance of anakinra decreased by 70% in severe renal insufficiency and by 75% in end stage renal disease (CrCl <30 ml/min); every other day dosing is recommended. Less than 2.5% of an administered dose was removed by dialysis (HD or CAPD); every other day dosing is recommended.*
- Baricitinib** *Approximately 75% of an administered dose was eliminated in the urine through filtration and active secretion, predominately as unchanged drug (69%). Renal function significantly affects baricitinib exposure. For patients with eGFR 30-60 ml/min, a dose of 2 mg once daily is recommended in the European product label and dose of 1 mg once daily is recommended in the US product label. Given short duration of therapy in COVID-19, 2 mg once daily advised for this indication. Not recommended for use in patients with eGFR <30 ml/min. Baricitinib is likely to be removed during CRRT; dose as in eGFR 30-60 ml/min.*
- Budesonide (inhaled)** *The metabolites of budesonide are excreted as such or in conjugated form mainly via the kidneys. No unchanged budesonide has been detected in the urine. There are no data regarding the specific use of inhaled budesonide in patients with renal impairment. No dose adjustment is required in renal impairment. Budesonide is unlikely to be dialysed.*
- Canakinumab** *Canakinumab is eliminated via intracellular catabolism. Due to its molecular size, little canakinumab is expected to be filtered by the kidney. No formal studies have been conducted to examine the pharmacokinetics of canakinumab administered subcutaneously in patients with renal impairment. Canakinumab is a human IgG immunoglobulin with large molecular size (~150 kDa), and little intact immunoglobulin is expected to be filtered by the kidney. Therefore, impaired renal function or renal replacement therapies are unlikely to affect the pharmacokinetics of canakinumab.*
- Dexamethasone (low dose)** *Dexamethasone is metabolised mainly in the liver, with up to 65% of the dose excreted unchanged in the urine. Particular care is required when considering the use of systemic corticosteroids in patients with renal insufficiency and frequent patient monitoring is necessary.*
- Fluvoxamine** *Fluvoxamine undergoes extensive metabolism in the liver. Fluvoxamine is excreted in the urine predominantly as metabolites, with approximately 2% of the administered dose recovered as unchanged drug. The mean minimum plasma concentrations in renally impaired patients (creatinine clearance 5-45 mL/min) after 4 and 6 weeks of treatment (50 mg twice daily, n=13) were comparable to each other, suggesting no accumulation of fluvoxamine in these patients. Patients with renal insufficiency (<10 ml/min) should start on a low dose and be carefully monitored.*
- Hydrocortisone** *Hydrocortisone is metabolised mainly in the liver and is minimally excreted in the urine. Particular care is required when using systemic hydrocortisone in patients with renal insufficiency (eGFR <50 ml/min); patient monitoring is advised. Hydrocortisone is unlikely to be dialysed.*

COVID-19 Therapies - Dose Recommendations for Patients with Renal Impairment

Charts updated 18 March 2022

Page 6 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

Notes – Host-directed Therapies (Licensed or Under Clinical Investigation) [continued]

- Infliximab** *The elimination pathways for infliximab have not been characterised but it is likely to be eliminated via its target antigen. Unchanged infliximab was not detected in urine.*
Infliximab has not been studied in patients with renal impairment. No dose recommendations can be made. Infliximab is not dialysed.
- Methylprednisolone** *Methylprednisolone is metabolized by CYP3A4 to inactive metabolites which are excreted in the urine.*
Particular care is required when considering the use of systemic corticosteroids in patients with renal insufficiency and frequent patient monitoring is necessary. Methylprednisolone is dialysed - dose as in normal renal function.
- Ruxolitinib** *Approximately 74% of an administered dose was eliminated in the urine, mainly as metabolites (<1% unchanged drug). Clearance of ruxolitinib metabolites decreases with increasing severity of renal impairment. The safety of increased exposure to these metabolites is unknown; close patient monitoring is advised in addition to dose adjustment recommendations.*
The US product label for ruxolitinib recommends to avoid in moderate/severe renal impairment if platelets <100. Avoid if eGFR <15 ml/min. Use with caution in patients on RRT. Administer post dialysis on dialysis days only. Ruxolitinib metabolites appeared to be dialysable to varying degrees by a 4-hour haemodialysis procedure. No data is available for dosing for patients on PD or CVVH.
- Sarilumab** *Sarilumab is not metabolised or excreted by the kidneys.*
No effect of renal impairment is expected.
Sarilumab's large molecular weight prevents clearance via glomerular filtration or RRT.
- Tocilizumab** *Tocilizumab is not metabolised or excreted by the kidneys.*
No effect of renal impairment is expected.
Tocilizumab's large molecular weight prevents clearance via glomerular filtration or RRT.

COVID-19 Therapies - Dose Recommendations for Patients with Renal Impairment

Charts updated 18 March 2022

Page 7 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

References

Antiviral Therapies

1. Zithromax (azithromycin) [Summary of Product Characteristics](#), Pfizer Ltd.
2. Zithromax (azithromycin) [Prescribing Information](#), Pfizer Inc.
3. Azithromycin: an assessment of its pharmacokinetics and therapeutic potential in CAPD. Kent JR, Almond MK, Dhillon S. *Perit Dial Int.* 2001;21(4):372-377.
4. The Renal Drug Handbook: The Ultimate Prescribing Guide for Renal Practitioners (5th Edition). Ashley C & Dunleavy A. CRC Press, Taylor & Francis Group, 2019.
5. Bamlanivimab and etesevimab [FDA Emergency Use Authorization](#), Lilly USA.
6. Ronapreve (casirivimab/imdevimab) [Summary of Product Characteristics](#), Roche Products Ltd.
7. REGEN-COV (casirivimab/imdevimab) [FDA Emergency Use Authorization](#), Regeneron Pharmaceuticals Inc.
8. Avigan (favipiravir) [Japanese Product Label](#), Toyama Chemical Co Ltd.
9. Lagevrio (molnupiravir) [Summary of Product Characteristics](#), Merck Sharp and Dohme Ltd.
10. Molnupiravir [FDA Emergency Use Authorization](#), Merck & Co Inc.
11. Yomesan (niclosamide) [Prescribing Information](#), Bayer (discontinued).
12. Niclosamide, [JoDrugs Health Inc.](#)
13. Paxlovid (nirmatrelvir/ritonavir) [Summary of Product Characteristics](#), Pfizer Ltd.
14. Paxlovid (nirmatrelvir/ritonavir) [FDA Emergency Use Authorization](#), Pfizer.
15. Alinia (nitazoxanide) [US Prescribing Information](#), Romark Pharmaceuticals.
16. Veklury (remdesivir) [Summary of Product Characteristics](#), Gilead Sciences Ltd.
17. Veklury (remdesivir) [US Prescribing Information](#), Gilead Sciences Inc.
18. Xevudy (sotrovimab) [Summary of Product Characteristics](#), GlaxoSmithKline.
19. Sotrovimab [FDA Emergency Use Authorization](#), GlaxoSmithKline.
20. Evusheld (tixagevimab/cilgavimab) [FDA Emergency Use Authorization](#), AstraZeneca.

Host-directed Therapies

21. Kineret (anakinra) [Summary of Product Characteristics](#), Swedish Orphan Biovitrum.
22. Kineret (anakinra) [US Prescribing Information](#), Swedish Orphan Biovitrum.
23. Olumiant (baracitinib) [Summary of Product Characteristics](#), Lilly.
24. Olumiant (baracitinib) [US Prescribing Information](#), Lilly USA.
25. Pulmicort (budesonide) [Summary of Product Characteristics](#), Astra Zeneca.
26. Pulmicort (budesonide) [US Prescribing Information](#), Astra Zeneca.
27. Ilaris (canakinumab) [US Prescribing Information](#), Novartis Pharmaceuticals.
28. Ilaris (canakinumab) [CHMP Assessment Report](#), European Medicines Agency.
29. Dexamethasone [Summary of Product Characteristics](#), Aspen.
30. Faverin (fluvoxamine) [Summary of Product Characteristics](#), Mylan.
31. Luvox (fluvoxamine) [US Prescribing Information](#), ANI Pharmaceuticals Inc.
32. Solu-Cortef (hydrocortisone) [Summary of Product Characteristics](#), Pfizer Ltd
33. Remicade(infliximab) [Summary of Product Characteristics](#), Merck Sharp and Dohme Ltd.
34. Remicade (infliximab) [US Prescribing Information](#), Janssen Pharmaceuticals.
35. Medrone (methylprednisolone) [Summary of Product Characteristics](#), Pfizer Ltd.
36. Medrol (methylprednisolone) [US Prescribing Information](#), Pfizer.
37. Jakavi (ruxolitinib) [Summary of Product Characteristics](#), Novartis Pharmaceuticals Ltd.
38. Jakafi (ruxolitinib) [US Prescribing Information](#), Incyte Corporation.
39. Pharmacokinetics and pharmacodynamics of orally administered ruxolitinib (INCB018424 phosphate) in renal and hepatic impairment patients. Chen X, Shi JG, Emm T, *et al.* *Clin Pharmacol Drug Dev.* 2014;3(1):34-42.
40. Kevzara (sarilumab) [Summary of Product Characteristics](#), Sanofi Genzyme.
41. RoActemra (tocilizumab, for infusion) [Summary of Product Characteristics](#), Roche Products Ltd.
42. Actemra (tocilizumab) [US Prescribing Information](#), Genentech Inc.